

THE TRANSFER OF PROPERTY (β) OF ROLEWICZ BY A UNIFORM QUOTIENT MAP

S. J. DILWORTH, DENKA KUTZAROVA, AND N. LOVASOA RANDRIANARIVONY

ABSTRACT. We provide a Laakso construction to prove that the property of having an equivalent norm with the property (β) of Rolewicz is qualitatively preserved via surjective uniform quotient mappings between separable Banach spaces. On the other hand, we show that the (β) -modulus is not quantitatively preserved via such a map by exhibiting two uniformly homeomorphic Banach spaces that do not have (β) -moduli of the same power-type even under renorming.

1. INTRODUCTION

A map $f : X \rightarrow Y$ between two metric spaces is called a uniform quotient mapping [5, Chapter 11] if there exist two nondecreasing maps $\omega, \Omega : [0, \infty) \rightarrow [0, \infty)$ such that $\omega(r) > 0$ for $r > 0$, and $\Omega(r) \rightarrow 0$ as $r \rightarrow 0$, and for every $x \in X$ and every $r > 0$,

$$B(f(x), \omega(r)) \subseteq f(B(x, r)) \subseteq B(f(x), \Omega(r)).$$

(In this article, all balls will be considered closed unless otherwise specified.) If there are constants $c, L > 0$ such that one can use $\omega(r) = cr$ and $\Omega(r) = Lr$, then the map f is called a Lipschitz quotient mapping. If the map f is surjective, the metric space Y is called a uniform quotient (resp. Lipschitz quotient) of the metric space X .

Closely related to the above definitions are the notions of coarse Lipschitz maps and coarse co-Lipschitz maps. A map $f : X \rightarrow Y$ between two metric spaces is called Lipschitz for large distances, or coarse Lipschitz, if for every $\delta > 0$ there exists $L(\delta) > 0$ such that $d_Y(f(x), f(x')) \leq L(\delta) d_X(x, x')$ for every $x, x' \in X$ with $d_X(x, x') \geq \delta$. The map f is called co-Lipschitz for large distances, or coarse co-Lipschitz, if for every $\delta > 0$ there exists $c(\delta) > 0$ such that for every $R \geq \delta$, for every $y, y' \in Y$ with $d_Y(y, y') \leq c(\delta)R$, for every $x \in f^{-1}(y)$, there exists $x' \in f^{-1}(y')$ with $d_X(x, x') \leq R$. It is known that when the metric spaces X and Y are metrically convex (for example if X and Y are Banach spaces), then a surjective uniform quotient mapping between X and Y is always coarse Lipschitz and coarse co-Lipschitz (see [5, Chapter 11] and [23, Section 3] for example).

2010 *Mathematics Subject Classification.* 46B80, 46B20, 46T99, 51F99.

Key words and phrases. Property (β) of Rolewicz, uniform quotient, Lipschitz quotient, Laakso construction.

The first author was partially supported by NSF grant DMS-1101490. The third author was partially supported by NSF grant DMS-1301591.

In [23] and in [9] the geometric property (β) of Rolewicz is applied to the study of the rigidity of Banach spaces under uniform quotient mappings. Rolewicz introduced property (β) as an isometric generalization of uniform convexity in the sense that a uniformly convex Banach space necessarily has property (β) (see [29] and [30]). Furthermore a result of Baudier, Kalton, and Lancien [4] implies that a separable reflexive Banach space X can be renormed to have property (β) if and only if the infinitely branching infinitely deep tree with its hyperbolic metric cannot bi-Lipschitz embed into X . Contrasted with Bourgain's metric characterization of superreflexivity (see [6] and [3]), this indicates that for separable spaces the isomorphic class of all Banach spaces with property (β) is also larger than the isomorphic class of all uniformly convex Banach spaces. We remark in passing that this statement is actually known to be true, see for example [18].

For the purpose of this article, the equivalent definition of property (β) given in [20] is used. A Banach space X has property (β) if for any $t \in (0, a]$, where the number $1 \leq a \leq 2$ depends on the space X , there exists $\delta > 0$ such that for any element x and sequence $(x_n)_{n \geq 1}$ in the unit ball B_X , if $\text{sep}((x_n)_{n \geq 1}) = \inf\{\|x_n - x_m\| : n \neq m; n, m \geq 1\} \geq t$, then there exists $n \geq 1$ satisfying

$$\frac{\|x - x_n\|}{2} \leq 1 - \delta.$$

The (β) -modulus was defined in [1] (although with a different notation) as

$$\bar{\beta}_X(t) = 1 - \sup \left\{ \inf \left\{ \frac{\|x - x_n\|}{2} : n \geq 1 \right\} : (x_n)_{n \geq 1} \subseteq B_X, x \in B_X, \text{sep}((x_n)_{n \geq 1}) \geq t \right\}.$$

In [9] it is proven that if a Banach space Y is a uniform quotient of another Banach space X with property (β) , then one can compare the (β) -modulus of X with the modulus of another asymptotic geometry on Y , namely asymptotic uniform smoothness, and under additional assumptions, asymptotic uniform convexity. (See [25] and [15] for the notion of asymptotic uniform smoothness and asymptotic uniform convexity. See also Section 4.) On the other hand, the authors in [24] use the (local) geometric property called Markov convexity to study the rigidity of Banach spaces under Lipschitz quotient mappings. They do this by comparing the Markov convexity of a space Y with the Markov convexity of a space X when Y is a Lipschitz quotient of a subset of X .

In light of these two approaches, the present article studies the asymptotic property (β) of Rolewicz on a Banach space Y that is a uniform quotient of another Banach space X with property (β) . This is done both in a qualitative sense and a quantitative sense. We show that the property of having an equivalent norm with property (β) is qualitatively preserved under a surjective uniform quotient mapping between separable Banach spaces. We do this by first assuming reflexivity on the range space in Section 2. In Section 3, we provide a modification of this approach to remove the reflexivity assumption.

In contrast, we show in Section 4 that quantitatively the (β) -modulus is not preserved under a surjective uniform quotient mapping. We also show that in full generality we cannot compare the (β) -modulus of the domain space with the modulus of asymptotic uniform convexity of the range space of a surjective uniform quotient map.

2. CASE WHEN THE RANGE SPACE IS REFLEXIVE

One of the main ingredients in this section is a metric characterization of being isomorphic to a space with property (β) . By combining the results of [9, Theorem 6.3] and [4, Theorem 1.2], we know that a separable reflexive Banach space Y has an equivalent norm with property (β) if and only if it does not contain any bi-Lipschitz copy of the metric tree \mathbb{T}_∞ of all finite subsets of \mathbb{N} with the shortest path metric.

Theorem 2.0.1. *Let X be a Banach space with (an equivalent norm with) property (β) . Let Y be a separable Banach space that is a uniform quotient of X . If Y is reflexive, then Y has an equivalent norm with property (β) .*

Proof. Let $f : X \rightarrow Y$ be a surjective uniform quotient mapping. Then f is coarse Lipschitz and coarse co-Lipschitz. For a contradiction, assume that Y does not admit any equivalent norm with property (β) . Denote by Σ a subset of Y that is bi-Lipschitz equivalent to \mathbb{T}_∞ , with a bi-Lipschitz equivalence denoted by $j : \Sigma \rightarrow \mathbb{T}_\infty$. The restriction $f|_{f^{-1}(\Sigma)} : f^{-1}(\Sigma) \rightarrow \Sigma$ is still a uniform quotient map that is coarse Lipschitz and coarse co-Lipschitz. As a result, since \mathbb{T}_∞ is discrete, the composition $j \circ f|_{f^{-1}(\Sigma)} : f^{-1}(\Sigma) \rightarrow \mathbb{T}_\infty$ is a (surjective) Lipschitz quotient map. The theorem then follows from the next proposition. \square

Proposition 2.0.2. *It is not possible that \mathbb{T}_∞ be a Lipschitz quotient of a subset of a Banach space with property (β) .*

Proof. We define a graph \mathbb{M}_∞ considered as a metric space, and we define a map $\phi : \mathbb{T}_\infty \rightarrow \mathbb{M}_\infty$. This map ϕ , although not a Lipschitz quotient map, will have a property still strong enough to draw a contradiction from the composition $\lambda = \phi \circ j \circ f|_{f^{-1}(\Sigma)} : f^{-1}(\Sigma) \rightarrow \mathbb{M}_\infty$, where $j \circ f|_{f^{-1}(\Sigma)} : f^{-1}(\Sigma) \rightarrow \mathbb{T}_\infty$ is a Lipschitz quotient map from a subset $f^{-1}(\Sigma)$ of a Banach space with property (β) . We support all these claims in the upcoming subsections. \square

2.1. Definition of the graph \mathbb{M}_∞ .

The graph \mathbb{M}_∞ is defined after a variant of the Laakso construction (see for example [8, 22, 24]). We define a sequence of graphs $(G_n)_{n \geq 1}$ such that $G_n \subseteq G_{n+1}$ for every $n \geq 1$ as follows. The graph G_1 is the following ordered graph. We start with a root. The root has one immediate descendant. This first descendant has countably infinitely many immediate descendants. Each one of those second generation vertices has only one immediate descendant, which is a common immediate descendant to all of them. The graph G_1 has diameter 3.



FIGURE 1. The graph G_1 . The “...” represents the fact that G_1 has infinitely many vertices at level 2.

To construct G_{n+1} ($n \geq 1$), we take a copy of G_1 and we rescale it so each edge has length the diameter of G_n . Next we replace each rescaled edge with a copy of G_n , matching the parent vertex of the rescaled edge with the oldest ancestor in the copy of G_n , and the child vertex of the rescaled edge with the youngest vertex in the copy of G_n . This gives all the vertices of G_{n+1} . Next we order the vertices of G_{n+1} by going downward from the root all the way to the youngest generation, which is now at distance 3^{n+1} from the root. Finally, we identify the root and the first 3^n generations of G_{n+1} with (the root and the first 3^n generations of) G_n . This way we have $G_n \subseteq G_{n+1}$.




FIGURE 2. The graph G_3 with the subsets G_1 and G_2 highlighted.

The graph \mathbb{M}_∞ is defined to be the union $\bigcup_{n=1}^{\infty} G_n$, considered as a metric space with the shortest path metric. The generational order of a given element of \mathbb{M}_∞ will be called its *level*. Note that the level of an element is also the length of a shortest path from the root to that element. We say that an element ν is a descendant of

another element μ , or that μ is an ancestor of ν , both denoted $\mu < \nu$, if there is a shortest path from the root to ν that passes through μ .

An element of \mathbb{M}_∞ will be called *non-branching* if it only has one immediate descendant; and it will be called *branching* if it has infinitely many immediate descendants. For each branching element of \mathbb{M}_∞ , we give a fraternal order among its immediate descendants by fixing a bijection between these descendants and the set \mathbb{N} of the natural numbers.

Remark 2.1.1. Laakso constructions, introduced in [21], have appeared elsewhere as a tool to investigate the geometry of metric spaces, including that of Banach spaces. The following list is not exhaustive: [31], [16], [24], [8], and recently [27]. We would like to point out the difference between our Laakso construction and that of [24, Section 3] and [8, Example 1.2] for example. First, our basic block G_1 has diameter three instead of four. Second, in the construction of our basic block G_1 , the first generation vertex has infinitely many immediate descendants instead of just two as in the standard Laakso construction. Third, when constructing G_{n+1} from G_n , we do *not* rescale down the copies of G_n . In our case, G_{n+1} has diameter three times as large as the diameter of G_n . We also note that even if the Laakso construction in [22] can be made so that the basic block has diameter three, our construction is still different from the construction in [22].

On the other hand we would like to mention that at the inception of this paper, our Laakso construction did have a basic block with diameter four. The proof went all the way through. However, as we prepared the final version of this article we realized that we did not need the last edge of the basic block, which brought us to the construction of \mathbb{M}_∞ as it is presented here.

2.2. Definition and properties of the map $\phi : \mathbb{T}_\infty \longrightarrow \mathbb{M}_\infty$.

We define $\phi(\text{root}) = \text{root}$. By induction, if $\phi(\bar{n})$ is already defined for an element \bar{n} of \mathbb{T}_∞ , and if \bar{m} is an immediate descendant of \bar{n} in \mathbb{T}_∞ , then

- set $\phi(\bar{m})$ to be the one immediate descendant of $\phi(\bar{n})$ if $\phi(\bar{n})$ is non-branching;
- otherwise, set $\phi(\bar{m})$ to be the immediate descendant of $\phi(\bar{n})$ in the same fraternal order among all immediate descendants of $\phi(\bar{n})$ as \bar{m} is among all immediate descendants of \bar{n} .

Before we list the properties of the map ϕ , let us make the following definitions.

Definition 2.2.1. Let $f : X \rightarrow Y$ be a map between two metric spaces X and Y . Let \mathcal{Q} be a subset of $X \times X$ and \mathcal{R} a subset of $Y \times Y$. We can, and will, think of \mathcal{Q} and \mathcal{R} as relations on X and Y respectively. We say that the map $f : X \rightarrow Y$ is

- \mathcal{Q} -Lipschitz if there exists $L^\mathcal{Q} > 0$ such that for every $x, x' \in X$, if $x \mathcal{Q} x'$ then $d_Y(f(x), f(x')) \leq L^\mathcal{Q} d_X(x, x')$;
- \mathcal{Q} -Lipschitz for large distances, or \mathcal{Q} -coarse Lipschitz, if for every $\delta > 0$ there exists $L^\mathcal{Q}(\delta) > 0$ such that $d_Y(f(x), f(x')) \leq L^\mathcal{Q}(\delta) d_X(x, x')$ for every $x, x' \in X$ with $x \mathcal{Q} x'$ and with $d_X(x, x') \geq \delta$;
- \mathcal{R} -co-Lipschitz if there exists $c^\mathcal{R} > 0$ such that for every $y, y' \in Y$ with $y \mathcal{R} y'$ and with $d_Y(y, y') \leq c^\mathcal{R} R$, for every $x \in f^{-1}(y)$, there exists $x' \in f^{-1}(y')$ with $d_X(x, x') \leq R$;
- \mathcal{R} -co-Lipschitz for large distances, or \mathcal{R} -coarse co-Lipschitz, if for every $\delta > 0$ there exists $c^\mathcal{R}(\delta) > 0$ such that for every $R \geq \delta$, for every $y, y' \in Y$

with $y\mathcal{R}y'$ and with $d_Y(y, y') \leq c^{\mathcal{R}}(\delta) R$, for every $x \in f^{-1}(y)$, there exists $x' \in f^{-1}(y')$ with $d_X(x, x') \leq R$.

In this article, the main relation we consider is that of ancestor-to-descendant. We now list the properties of the map ϕ in the following lemma.

Lemma 2.2.2.

- (1) ϕ is surjective, and preserves levels.
- (2) ϕ is 1-Lipschitz.
- (3) ϕ is ancestor-to-descendant 1-co-Lipschitz in the following sense:

$\forall \mu, \nu \in \mathbb{M}_\infty$ with $\mu < \nu$, $\forall \bar{n} \in \phi^{-1}(\mu)$, $\exists \bar{m} \in \phi^{-1}(\nu)$ such that $d_{\mathbb{T}_\infty}(\bar{n}, \bar{m}) = d_{\mathbb{M}_\infty}(\mu, \nu)$.

Proof.

- (1) This claim is easy.
- (2) Let $\bar{n}, \bar{m} \in \mathbb{T}_\infty$. We consider two cases.
 - Case $\bar{n} < \bar{m}$:

$$\begin{aligned} d_{\mathbb{M}_\infty}(\phi(\bar{n}), \phi(\bar{m})) &= \text{level}(\phi(\bar{m})) - \text{level}(\phi(\bar{n})) \\ &= \text{level}(\bar{m}) - \text{level}(\bar{n}) \\ &= d_{\mathbb{T}_\infty}(\bar{n}, \bar{m}). \end{aligned}$$

- Case \bar{n} and \bar{m} are not comparable:
Let $\bar{o} \in \mathbb{T}_\infty$ be the closest common ancestor to \bar{n} and \bar{m} in \mathbb{T}_∞ ; and let $\mu \in \mathbb{M}_\infty$ be the closest common ancestor to $\phi(\bar{n})$ and $\phi(\bar{m})$ in \mathbb{M}_∞ . Since $\phi(\bar{o})$ is also a common ancestor to $\phi(\bar{n})$ and $\phi(\bar{m})$, the definition of μ gives us that $\text{level}(\phi(\bar{o})) \leq \text{level}(\mu)$. Hence we have:

$$\begin{aligned} d_{\mathbb{M}_\infty}(\phi(\bar{n}), \phi(\bar{m})) &\leq d_{\mathbb{M}_\infty}(\phi(\bar{n}), \mu) + d_{\mathbb{M}_\infty}(\mu, \phi(\bar{m})) \\ &= \text{level}(\phi(\bar{n})) - \text{level}(\mu) + \text{level}(\phi(\bar{m})) - \text{level}(\mu) \\ &\leq \text{level}(\phi(\bar{n})) - \text{level}(\phi(\bar{o})) + \text{level}(\phi(\bar{m})) - \text{level}(\phi(\bar{o})) \\ &= \text{level}(\bar{n}) - \text{level}(\bar{o}) + \text{level}(\bar{m}) - \text{level}(\bar{o}) \\ &= d_{\mathbb{T}_\infty}(\bar{n}, \bar{m}). \end{aligned}$$

- (3) Let $\mu < \nu \in \mathbb{M}_\infty$, and let $\bar{n} \in \phi^{-1}(\mu)$. Choose a path in \mathbb{M}_∞ from μ down to ν . Record (every time such is defined) all the fraternity orders of all elements along that path. Then, starting from μ , lift that path node by node, inductively on the level of the node, as follows:
 - $\phi(\bar{n}) = \mu$.
 - Assume all nodes on the path at distance less than or equal to k from μ have been lifted. Let ζ be on the path, at distance $k+1$ from μ . Let ζ' be the immediate ancestor of ζ (which is well-defined since we are

following a given path), and let $\bar{n}' \in \phi^{-1}(\zeta')$ be the lift of ζ' chosen according to the induction hypothesis. We consider two cases.

- * If ζ' is a non-branching node, then lift ζ to any immediate descendant of \bar{n}' .
- * If ζ' is a branching node, then lift ζ to the immediate descendant of \bar{n}' of the same fraternal order in \mathbb{T}_∞ as ζ is among all immediate descendants of ζ' .
- Doing this way, we eventually get to an element \bar{m} of \mathbb{T}_∞ such that $\phi(\bar{m}) = \nu$ with $\bar{m} > \bar{n}$ and $d_{\mathbb{M}_\infty}(\mu, \nu) = d_{\mathbb{T}_\infty}(\bar{n}, \bar{m})$.

□

2.3. Fork argument.

The fork argument was introduced in [23, proof of Theorem 4.1] for maps that are coarse co-Lipschitz. Here we reduce its hypothesis when we apply it to a map λ that is only ancestor-to-descendant coarse co-Lipschitz.

First let us fix some notation. Consider a metric space \mathcal{S} and a surjective map $\lambda : \mathcal{S} \rightarrow \mathbb{M}_\infty$ that is ancestor-to-descendant coarse co-Lipschitz. For $\delta > 0$, consider the (nonempty) set $\mathcal{C}(\delta)$ of all $c > 0$ such that for all $R \geq \delta$, for all $\sigma \in \mathcal{S}$, and for all $\nu > \lambda(\sigma)$ in \mathbb{M}_∞ with $d_{\mathbb{M}_\infty}(\lambda(\sigma), \nu) < cR$, there exists $\sigma' \in \mathcal{S}$ with $d_{\mathcal{S}}(\sigma, \sigma') \leq R$ and $\lambda(\sigma') = \nu$. Denote by c_δ^{ATD} the supremum of the set $\mathcal{C}(\delta)$. It is left to the reader to check (see also [23, Lemma 3.3]) that c_δ^{ATD} is attained, and that the family $(c_\delta^{\text{ATD}})_{\delta > 0}$ is increasing. Denote by $c_\infty^{\text{ATD}} := \sup \{c_\delta^{\text{ATD}} : \delta > 0\} = \lim_{\delta \rightarrow \infty} c_\delta^{\text{ATD}}$.

With these notation, we present the fork argument for ancestor-to-descendant coarse co-Lipschitz maps.

Proposition 2.3.1. *For a metric space \mathcal{S} , let $\lambda : \mathcal{S} \rightarrow \mathbb{M}_\infty$ be a surjective map that is ancestor-to-descendant co-Lipschitz for large distances. Assume that the quantity c_∞^{ATD} for the map λ is finite. Then for every $\varepsilon > 0$, one can find $r > 0$ as large as we want, as well as elements μ_0, μ_1 , and $(\mu_{2,k})_{k \geq 1}$ in \mathbb{M}_∞ , and elements σ_0, σ_1 , and $(\sigma_{2,k})_{k \geq 1}$ in \mathcal{S} so that $\lambda(\sigma_0) = \mu_0$, $\lambda(\sigma_1) = \mu_1$, and $\lambda(\sigma_{2,k}) = \mu_{2,k}$ for every $k \geq 1$; with the σ 's and the μ 's sitting in an (approximate) fork position as follows:*

$$d_{\mathbb{M}_\infty}(\mu_0, \mu_1) = d_{\mathbb{M}_\infty}(\mu_1, \mu_{2,k}) = \frac{d_{\mathbb{M}_\infty}(\mu_0, \mu_{2,k})}{2} = r;$$

and

$$\begin{aligned} d_{\mathcal{S}}(\sigma_0, \sigma_1) &\leq (1 + 3\varepsilon) \frac{r}{c_\infty^{\text{ATD}}}, \\ d_{\mathcal{S}}(\sigma_1, \sigma_{2,k}) &\leq (1 + 3\varepsilon) \frac{r}{c_\infty^{\text{ATD}}}, \\ \frac{d_{\mathcal{S}}(\sigma_0, \sigma_{2,k})}{2} &> (1 - 80\varepsilon) \frac{r}{c_\infty^{\text{ATD}}}. \end{aligned}$$

Proof. Let $\varepsilon > 0$. Let $\delta_0 = \frac{3^{n_0}}{c_\infty^{\text{ATD}}}$ for some large enough n_0 such that $c_\infty^{\text{ATD}} < (1 + \varepsilon)c_{\delta_0}^{\text{ATD}}$. Then for any $\delta \geq \delta_0$, we have

$$c_{\delta_0}^{\text{ATD}} \leq c_\delta^{\text{ATD}} \leq c_\infty^{\text{ATD}} < (1 + \varepsilon)c_{\delta_0}^{\text{ATD}} \leq (1 + \varepsilon)c_\delta^{\text{ATD}} \leq (1 + \varepsilon)c_\infty^{\text{ATD}}.$$

Since c_δ^{ATD} is a maximum, since $(1 + \varepsilon)c_\delta^{\text{ATD}} > c_\delta^{\text{ATD}}$, and because of the ancestor-to-descendant coarse co-Lipschitz condition on λ , we can find a number $R \geq \delta$, and

elements $\sigma \in \mathcal{S}$ and $\nu > \lambda(\sigma) \in \mathbb{M}_\infty$ with $c_\delta^{\text{ATD}} R \leq d_{\mathbb{M}_\infty}(\nu, \lambda(\sigma)) < (1 + \varepsilon) c_\delta^{\text{ATD}} R$ such that

$$\forall \sigma' \in \mathcal{S}, (\lambda(\sigma') = \nu) \implies (d_{\mathcal{S}}(\sigma, \sigma') > R).$$

In passing, note that we can then make our choice of δ_0 in such a way that $d_{\mathbb{M}_\infty}(\nu, \lambda(\sigma))$ is as large as we want.

Let us call $\mu := \lambda(\sigma)$. Let $N \in \mathbb{N}$ be such that $3^N \leq \frac{d_{\mathbb{M}_\infty}(\mu, \nu)}{2} < 3^{N+1}$. Since ν is a descendant of μ , we have $d_{\mathbb{M}_\infty}(\mu, \nu) = \text{level}(\nu) - \text{level}(\mu)$, and hence there must exist an integer $p \geq 0$ such that $\text{level}(\mu) \leq p3^N < (p+1)3^N \leq \text{level}(\nu)$. In fact, one can take p to be the largest integer such that $(p-1)3^N < \text{level}(\mu)$. Furthermore, for $\delta \geq 81\delta_0$, we must necessarily have $3^{n_0} < 3^{N-2}$. In fact, $\delta \geq 54(1 + \varepsilon)\delta_0$ if ε is small enough. Then since $(1 + \varepsilon) \geq c_\infty^{\text{ATD}}/c_\delta^{\text{ATD}}$, and since $c_\infty^{\text{ATD}}\delta_0 = 3^{n_0}$, we have

$$2 \cdot 3^{N+1} > d_{\mathbb{M}_\infty}(\mu, \nu) \geq c_\delta^{\text{ATD}} R \geq c_\delta^{\text{ATD}} \delta \geq 6 \cdot 9 \cdot 3^{n_0}.$$

We will now identify the elements μ_0, μ_1 and $\mu_{2,k} (k \geq 1)$ of \mathbb{M}_∞ that we need for our purpose. First, we fix a path \mathcal{P} downward from μ to ν . We consider the elements a and b of \mathcal{P} at levels $p \cdot 3^N$ and $(p+1)3^N$ respectively. Note that the segment $[a, b]$ has length 3^N . Divide $[a, b]$ into three equal-length subsegments, and denote by μ_0 the element of \mathcal{P} at level $p \cdot 3^N + 3^{N-1}$, and by μ_3 the one at level $p \cdot 3^N + 2 \cdot 3^{N-1}$. Note that $d_{\mathbb{M}_\infty}(\mu, \mu_0) \geq 3^{N-1}$ and $d_{\mathbb{M}_\infty}(\mu_3, \nu) \geq 3^{N-1}$. Divide the segment $[\mu_0, \mu_3]$ into three again, and consider the following vertices: the element μ_1 of \mathcal{P} at level $p \cdot 3^N + 3^{N-1} + 3^{N-2}$, and the elements $\mu_{2,k} (k \geq 1)$ of \mathbb{M}_∞ at level $p \cdot 3^N + 3^{N-1} + 2 \cdot 3^{N-2}$ that lie between μ_1 and the element μ_3 already previously mentioned. Note that although only one of the $\mu_{2,k}$'s belongs to the original path \mathcal{P} , the succession $\mu < \mu_0 < \mu_1 < \mu_{2,k} < \mu_3 < \nu$ can be extended to a shortest path from μ to ν for any $k \geq 1$.

Let us set

$$d_{\mathbb{M}_\infty}(\mu_1, \mu_0) = d_{\mathbb{M}_\infty}(\mu_{2,k}, \mu_1) = d_{\mathbb{M}_\infty}(\mu_3, \mu_{2,k}) := r (= 3^{N-2} \geq c_\infty^{\text{ATD}}\delta_0).$$

Then note that $d_{\mathbb{M}_\infty}(\mu_{2,k}, \nu) = d_{\mathbb{M}_\infty}(\mu, \nu) - d_{\mathbb{M}_\infty}(\mu, \mu_0) - 2r$ for each $k \geq 1$. We will now lift these elements one by one starting from μ using the ancestor-to-descendant coarse co-Lipschitz condition.

- (1) $d_{\mathbb{M}_\infty}(\mu, \mu_0) < c_{\delta_0}^{\text{ATD}} \cdot \frac{(1 + \varepsilon)}{c_{\delta_0}^{\text{ATD}}} d_{\mathbb{M}_\infty}(\mu, \mu_0)$; and $\frac{(1 + \varepsilon)}{c_{\delta_0}^{\text{ATD}}} d_{\mathbb{M}_\infty}(\mu, \mu_0) \geq \delta_0$ because $d_{\mathbb{M}_\infty}(\mu, \mu_0) \geq 3^{N-1} \geq c_\infty^{\text{ATD}}\delta_0 \geq c_{\delta_0}^{\text{ATD}}\delta_0$. So there exists an element $\sigma_0 \in \mathcal{S}$ with $d_{\mathcal{S}}(\sigma, \sigma_0) \leq \frac{1 + \varepsilon}{c_{\delta_0}^{\text{ATD}}} d_{\mathbb{M}_\infty}(\mu, \mu_0)$ such that $\lambda(\sigma_0) = \mu_0$.
- (2) $d_{\mathbb{M}_\infty}(\mu_0, \mu_1) = r < c_{\delta_0}^{\text{ATD}} \cdot \frac{1 + \varepsilon}{c_{\delta_0}^{\text{ATD}}} r$; and $\frac{1 + \varepsilon}{c_{\delta_0}^{\text{ATD}}} r \geq \delta_0$ since $r = 3^{N-2} \geq c_\infty^{\text{ATD}}\delta_0$. So there is an element $\sigma_1 \in \mathcal{S}$ with $d_{\mathcal{S}}(\sigma_0, \sigma_1) \leq \frac{1 + \varepsilon}{c_{\delta_0}^{\text{ATD}}} r$ such that $\lambda(\sigma_1) = \mu_1$.
- (3) Similarly, $d_{\mathbb{M}_\infty}(\mu_1, \mu_{2,k}) = r$ for each $k \geq 1$, so there is an element $\sigma_{2,k} \in \mathcal{S}$ with $d_{\mathcal{S}}(\sigma_1, \sigma_{2,k}) \leq \frac{1 + \varepsilon}{c_{\delta_0}^{\text{ATD}}} r$ such that $\lambda(\sigma_{2,k}) = \mu_{2,k}$.

- (4) Finally, $d_{\mathbb{M}_\infty}(\mu_{2,k}, \nu) = d_{\mathbb{M}_\infty}(\mu, \nu) - d_{\mathbb{M}_\infty}(\mu, \mu_0) - 2r < c_{\delta_0}^{\text{ATD}} \cdot \frac{1+\varepsilon}{c_{\delta_0}^{\text{ATD}}} [d_{\mathbb{M}_\infty}(\mu, \nu) - d_{\mathbb{M}_\infty}(\mu, \mu_0) - 2r]$; and $\frac{1+\varepsilon}{c_{\delta_0}^{\text{ATD}}} [d_{\mathbb{M}_\infty}(\mu, \nu) - d_{\mathbb{M}_\infty}(\mu, \mu_0) - 2r] \geq \frac{1+\varepsilon}{c_{\delta_0}^{\text{ATD}}} r \geq \delta_0$, so there exists an element $\sigma^{(k)} \in \mathcal{S}$ with $d_{\mathcal{S}}(\sigma_{2,k}, \sigma^{(k)}) \leq \frac{1+\varepsilon}{c_{\delta_0}^{\text{ATD}}} [d_{\mathbb{M}_\infty}(\mu, \nu) - d_{\mathbb{M}_\infty}(\mu, \mu_0) - 2r]$ such that $\lambda(\sigma^{(k)}) = \nu$.

We claim that as a result, we must have

$$d_{\mathcal{S}}(\sigma_{2,k}, \sigma_0) > [1 - (3^4 - 1)\varepsilon] \cdot \frac{2r}{c_{\delta_0}^{\text{ATD}}}.$$

In fact by the choice of μ and ν , we have $d_{\mathcal{S}}(\sigma, \sigma^{(k)}) > R$. So if $d_{\mathcal{S}}(\sigma_{2,k}, \sigma_0) \leq \alpha \cdot \frac{2r}{c_{\delta_0}^{\text{ATD}}}$, then

$$\begin{aligned} R &< d_{\mathcal{S}}(\sigma, \sigma^{(k)}) \\ &\leq d_{\mathcal{S}}(\sigma, \sigma_0) + d_{\mathcal{S}}(\sigma_0, \sigma_{2,k}) + d_{\mathcal{S}}(\sigma_{2,k}, \sigma^{(k)}) \\ &\leq \frac{1+\varepsilon}{c_{\delta_0}^{\text{ATD}}} d_{\mathbb{M}_\infty}(\mu, \mu_0) + \frac{\alpha}{c_{\delta_0}^{\text{ATD}}} 2r + \frac{1+\varepsilon}{c_{\delta_0}^{\text{ATD}}} (d_{\mathbb{M}_\infty}(\mu, \nu) - d_{\mathbb{M}_\infty}(\mu, \mu_0) - 2r) \\ &= \frac{\alpha}{c_{\delta_0}^{\text{ATD}}} 2r + \frac{1+\varepsilon}{c_{\delta_0}^{\text{ATD}}} (d_{\mathbb{M}_\infty}(\mu, \nu) - 2r) \\ &= \frac{d_{\mathbb{M}_\infty}(\mu, \nu)}{c_{\delta_0}^{\text{ATD}}} [t\alpha + (1-t)(1+\varepsilon)], \text{ where } t = \frac{2r}{d_{\mathbb{M}_\infty}(\mu, \nu)} \\ &< \frac{(1+\varepsilon)c_{\infty}^{\text{ATD}} R}{c_{\delta_0}^{\text{ATD}}} [t\alpha + (1-t)(1+\varepsilon)] \\ &< (1+\varepsilon)^2 [t\alpha + (1-t)(1+\varepsilon)] R. \end{aligned}$$

This would require that $\alpha > (1+\varepsilon) - \frac{1}{t} \left[(1+\varepsilon) - \frac{1}{(1+\varepsilon)^2} \right]$. But from the definition of N , we have $d_{\mathbb{M}_\infty}(\mu, \nu) < 2 \cdot 3^{N+1}$. So, since $r = 3^{N-2}$, we have

$$\frac{1}{t} = \frac{d_{\mathbb{M}_\infty}(\mu, \nu)}{2r} < 3^3,$$

and hence

$$\alpha > (1+\varepsilon) - 3^3 \left[(1+\varepsilon) - \frac{1}{(1+\varepsilon)^2} \right] \geq (1+\varepsilon) - 3^3(3\varepsilon).$$

To summarize, we have $\sigma_0 \in \lambda^{-1}(\mu_0)$, $\sigma_1 \in \lambda^{-1}(\mu_1)$, and $\sigma_{2,k} \in \lambda^{-1}(\mu_{2,k})$ ($k \geq 1$), and with ε small enough we have

$$d_{\mathcal{S}}(\sigma_0, \sigma_1) \leq (1 + \varepsilon) \frac{r}{c_{\delta_0}^{\text{ATD}}} < (1 + \varepsilon)^2 \frac{r}{c_{\infty}^{\text{ATD}}} \leq (1 + 3\varepsilon) \frac{r}{c_{\infty}^{\text{ATD}}},$$

and

$$d_{\mathcal{S}}(\sigma_1, \sigma_{2,k}) \leq (1 + \varepsilon) \frac{r}{c_{\delta_0}^{\text{ATD}}} < (1 + \varepsilon)^2 \frac{r}{c_{\infty}^{\text{ATD}}} \leq (1 + 3\varepsilon) \frac{r}{c_{\infty}^{\text{ATD}}},$$

but

$$d_{\mathcal{S}}(\sigma_0, \sigma_{2,k}) > (1 - 80\varepsilon) \frac{2r}{c_{\delta_0}^{\text{ATD}}} \geq (1 - 80\varepsilon) \frac{2r}{c_{\infty}^{\text{ATD}}}.$$

This finishes the proof. \square

2.4. Proof of Proposition 2.0.2.

We apply Proposition 2.3.1 to the set $\mathcal{S} := f^{-1}(\Sigma) = f^{-1}(j^{-1}(\mathbb{T}_{\infty}))$ and the map $\lambda = \phi \circ j \circ f|_{\mathcal{S}}$. Recall that this particular map is also Lipschitz, and hence the hypothesis of Proposition 2.3.1 applies since $c_{\delta}^{\text{ATD}} \leq \text{Lip}(\lambda)$ for all $\delta > 0$. We find the elements $\mu_0, \mu_1, (\mu_{2,k})_{k \geq 1} \in \mathbb{M}_{\infty}$, and $\sigma_0, \sigma_1, (\sigma_{2,k})_{k \geq 1} \in \mathcal{S} \subseteq X$ as given by Proposition 2.3.1. We have for all $k \geq 1$,

$$\|\sigma_0 - \sigma_1\|_X \leq (1 + 3\varepsilon) \frac{r}{c_{\infty}^{\text{ATD}}},$$

and

$$\|\sigma_1 - \sigma_{2,k}\|_X \leq (1 + 3\varepsilon) \frac{r}{c_{\infty}^{\text{ATD}}},$$

but

$$\|\sigma_0 - \sigma_{2,k}\|_X > (1 - 80\varepsilon) \frac{2r}{c_{\infty}^{\text{ATD}}}.$$

On the other hand, since λ is Lipschitz, we have for all $k \neq l$,

$$\|\sigma_{2,k} - \sigma_{2,l}\|_X \geq \frac{1}{\text{Lip}(\lambda)} d_{\mathbb{M}_{\infty}}(\mu_{2,k}, \mu_{2,l}) = \frac{1}{\text{Lip}(\lambda)} 2r.$$

Since X has property (β) , this implies that we must have for ε small enough,

$$\bar{\beta}_X \left(\frac{c_{\infty}^{\text{ATD}}}{\text{Lip}(\lambda)} \right) \leq \bar{\beta}_X \left(\frac{2c_{\infty}^{\text{ATD}}}{(1 + 3\varepsilon)\text{Lip}(\lambda)} \right) \leq \frac{83\varepsilon}{1 + 3\varepsilon}.$$

Since ε is arbitrary, we then get that

$$\bar{\beta}_X \left(\frac{c_{\infty}^{\text{ATD}}}{\text{Lip}(\lambda)} \right) = 0.$$

This contradiction finishes the proof. \square

3. REMOVING THE REFLEXIVITY ASSUMPTION

James proves many characterizations of reflexivity in [13]. We highlight in the following lemma the one that we need. For this purpose, denote by \mathbb{T} the set of all finite subsets of \mathbb{N} , considered in graph-theoretic terms as a tree. Note that this time \mathbb{T} is only considered as a tree to organize its elements, but we do not put any metric on \mathbb{T} . As a set, we have $\mathbb{T} = \mathbb{T}_\infty$.

Lemma 3.0.1 (James). *Let X be a non-reflexive Banach space. Then for every $0 < \theta < 1$, there exists a sequence $(u_n)_{n \geq 1} \subseteq B_X$ such that*

- (1) *the map from \mathbb{T} to X given by $\{n_1 < n_2 < \dots < n_k\} \mapsto u_{n_1} + \dots + u_{n_k}$ is one-to-one,*
- (2) *for every $n_1 < \dots < n_k$, we have*

$$\theta k \leq \|u_{n_1} + \dots + u_{n_k}\| \leq k,$$

- (3) *and for every $n_1 < \dots < n_k < m_1 < \dots < m_l$, we have*

$$\frac{\theta}{3}(k+l) \leq \|(u_{n_1} + \dots + u_{n_k}) - (u_{m_1} + \dots + u_{m_l})\| \leq k+l.$$

Proof. Let $\theta \in (0, 1)$. From item (31) in [13], we can find sequences $(u_n)_{n \geq 1} \subseteq B_X$ and $(u_n^*)_{n \geq 1} \subseteq B_{X^*}$ such that $u_n^*(u_k) = \theta$ if $n \leq k$, and $u_n^*(u_k) = 0$ if $n > k$.

- (1) Let $\{n_1 < \dots < n_k\} \neq \{m_1 < \dots < m_l\}$. Denote $\{n'_1 < \dots < n'_s\} = \{n_1 < \dots < n_k\} \Delta \{m_1 < \dots < m_l\}$, where $s \leq k+l$; and let signs $(\varepsilon_i)_{1 \leq i \leq s} \subseteq \{-1, 1\}$ be such that

$$(u_{n_1} + \dots + u_{n_k}) - (u_{m_1} + \dots + u_{m_l}) = \varepsilon_1 u_{n'_1} + \dots + \varepsilon_s u_{n'_s}.$$

Then we have

$$\|(u_{n_1} + \dots + u_{n_k}) - (u_{m_1} + \dots + u_{m_l})\| \geq |u_{n'_s}^* (\varepsilon_1 u_{n'_1} + \dots + \varepsilon_s u_{n'_s})| = |\varepsilon_s \theta| > 0.$$

- (2) For $n_1 < \dots < n_k$, we have:

$$\|u_{n_1} + \dots + u_{n_k}\| \geq u_{n_1}^* (u_{n_1} + \dots + u_{n_k}) = \theta k.$$

The other inequality follows from the triangle inequality.

- (3) Let $n_1 < \dots < n_k < m_1 < \dots < m_l$. On the one hand, we have

$$\|(u_{n_1} + \dots + u_{n_k}) - (u_{m_1} + \dots + u_{m_l})\| \geq |u_{m_1}^* ((u_{n_1} + \dots + u_{n_k}) - (u_{m_1} + \dots + u_{m_l}))| = \theta l,$$

and on the other hand,

$$\|(u_{n_1} + \dots + u_{n_k}) - (u_{m_1} + \dots + u_{m_l})\| \geq |u_{n_1}^* ((u_{n_1} + \dots + u_{n_k}) - (u_{m_1} + \dots + u_{m_l}))| = \theta |k-l|.$$

We get the left inequality by noting that $\max\{l, |k-l|\} \geq \frac{1}{3}(k+l)$. Again the inequality on the right follows from the triangle inequality.

□

We now state the main result in this section.

Theorem 3.0.2. *Assume that a Banach space Y is a uniform quotient of a Banach space X . If X has (an equivalent norm with) property (β) , then Y has to be reflexive.*

Proof. Let $f : X \rightarrow Y$ be a surjective uniform quotient map. Note that f is coarse Lipschitz and coarse co-Lipschitz. Assume that Y is not reflexive. We apply James' characterization with $\theta = 3/4$. For $J = \{n_1 < n_2 < \dots < n_k\} \in \mathbb{T}$, set $v_J := u_{n_1} + \dots + u_{n_k}$. We then get that if $\max J < \min J'$, then

$$(1) \quad \begin{cases} \frac{1}{4}|J| \leq \|v_J\|_Y \leq |J|, \\ \frac{1}{4}|J'| \leq \|v_{J'}\|_Y \leq |J'|, \\ \frac{1}{4}(|J| + |J'|) \leq \|v_J - v_{J'}\|_Y \leq (|J| + |J'|). \end{cases}$$

Consider the subset $\mathcal{T} := \{v_J \in Y : J \in \mathbb{T}\} \subseteq Y$, which is a metric space. Since the correspondence $J \mapsto v_J$ is one-to-one, the set \mathcal{T} is in one-to-one and onto correspondence with \mathbb{T} , and we extend the notions of levels, ancestors, and descendants to elements of \mathcal{T} as well. Consider the composition

$$\tilde{\phi} := (\phi \circ \equiv) : \mathcal{T} \equiv \mathbb{T}_\infty \rightarrow \mathbb{M}_\infty,$$

where ϕ is the same map as in Subsection 2.2, and \equiv is the natural bijection between \mathcal{T} and \mathbb{T}_∞ . The bijection $\mathcal{T} \equiv \mathbb{T}_\infty$ is ancestor-to-descendant bi-Lipschitz by James' characterization as expressed in (1) above. And hence the composition $\tilde{\phi} : \mathcal{T} \rightarrow \mathbb{M}_\infty$, considered as a map between metric spaces, is ancestor-to-descendant co-Lipschitz.

Next, consider the restriction $f|_{f^{-1}(\mathcal{T})} : f^{-1}(\mathcal{T}) \rightarrow \mathcal{T} \subseteq Y$, which is a surjective uniform quotient mapping that is coarse Lipschitz and coarse co-Lipschitz. As a result, the composition $\tilde{\lambda} := \tilde{\phi} \circ f|_{f^{-1}(\mathcal{T})} : f^{-1}(\mathcal{T}) \rightarrow \mathbb{M}_\infty$ is also ancestor-to-descendant co-Lipschitz. Note that $c_\infty^{\text{ATD}}(\tilde{\lambda})$ is finite since the inequalities (1) also give that the map $\tilde{\phi}$ is ancestor-to-descendant Lipschitz. We can thus apply the fork argument (Proposition 2.3.1) to $\tilde{\lambda}$, producing elements $\mu_0, \mu_1, \mu_{2,k}$ ($k \geq 1$) in \mathbb{M}_∞ with $d_{\mathbb{M}_\infty}(\mu_0, \mu_1) = d_{\mathbb{M}_\infty}(\mu_1, \mu_{2,k}) = d_{\mathbb{M}_\infty}(\mu_0, \mu_{2,k})/2 = r = 3^{N-2}$; as well as respective $\tilde{\lambda}$ -preimages $\sigma_0, \sigma_1, \sigma_{2,k}$ ($k \geq 1$) in $f^{-1}(\mathcal{T}) \subseteq X$ with

$$(2) \quad \begin{cases} \|\sigma_0 - \sigma_1\|_X, \|\sigma_1 - \sigma_{2,k}\|_X \leq (1 + 3\varepsilon) \frac{r}{c_\infty^{\text{ATD}}(\tilde{\lambda})} \\ \|\sigma_0 - \sigma_{2,k}\|_X > (1 - 80\varepsilon) \frac{2r}{c_\infty^{\text{ATD}}(\tilde{\lambda})}. \end{cases}$$

Let us denote $v^{(0)} := f(\sigma_0)$, $v^{(1)} := f(\sigma_1)$, and $v_{J^{(k)}} := f(\sigma_{2,k})$ (for all $k \geq 1$), which are elements of the subset \mathcal{T} of Y . Recall that the map ϕ preserves levels, so

$$\begin{cases} \text{level}(v^{(0)}) = \text{level}(\mu_0), \\ \text{level}(v^{(1)}) = \text{level}(\mu_1) = \text{level}(\mu_0) + 3^{N-2}, \\ \text{level}(v_{J^{(k)}}) = \text{level}(\mu_{2,k}) = \text{level}(\mu_1) + 3^{N-2}. \end{cases}$$

But $\text{level}(v_{J^{(k)}}) = |J^{(k)}|$, so $|J^{(k)}| = \text{level}(\mu_1) + 3^{N-2}$ for all $k \geq 1$.

Consider the set of all elements of \mathbb{T} with infinitely (not necessarily immediate) descendants among the $J^{(k)}$'s. This set is nonempty since the root \emptyset belongs to it. Now let J be an element of this set such that $\text{level}(J) = |J|$ is maximal. Let $(J^{(k_n)})_{n \geq 1}$ be the infinitely many descendants of J among the $J^{(k)}$'s. Write $J^{(k_n)} =: J \cup I^{(n)}$ with $\max J < \min I^{(n)}$. Then for every $n \geq 1$, we must have

$$(3) \quad |I^{(n)}| \geq 3^{N-2}.$$

In fact, the map $\phi : \mathbb{T}_\infty \rightarrow \mathbb{M}_\infty$ is 1-Lipschitz, so for any $k \neq j$ we have

$$2 \cdot 3^{N-2} = d_{\mathbb{M}_\infty}(\mu_{2,k}, \mu_{2,j}) \leq d_{\mathbb{T}_\infty}(J^{(k)}, J^{(j)}),$$

i.e. the closest common ancestor to $J^{(k)}$ and $J^{(j)}$ is at least 3^{N-2} generations back.

Next, consider the immediate descendants $(J \cup \{m\})_{m > \max(J)}$ of J . By the fact that $\text{level}(J)$ is maximal among the elements of \mathbb{T} with infinitely many descendants among the $J^{(k)}$'s, we must find an infinite sequence $m_1 < m_2 < \dots$ such that

- each $J \cup \{m_j\}$ has descendants among the $J^{(k_n)}$'s,
- and each such $J \cup \{m_j\}$ has only finitely many descendants among the $J^{(k_n)}$'s.

Take a further subsequence of $(m_j)_{j \geq 1}$, still denoted $(m_j)_{j \geq 1}$, such that if we denote by $J^{(k_{n_j})} = J \cup I^{(n_j)}$ the descendant of $J \cup \{m_j\}$ with $\max(I^{(n_j)})$ being maximal, then $\max(I^{(n_j)}) < m_{j+1} = \min(I^{(n_{j+1})})$.

Then, by James, we have

$$\|v_{J^{(k_{n_j})}} - v_{J^{(k_{n_{j'}})}}\|_Y \geq \frac{1}{4} (|I^{(n_j)}| + |I^{(n_{j'})}|) \geq \frac{1}{2} \cdot 3^{N-2}.$$

(The last inequality comes from inequality (3).) Using the Lipschitz condition for large distances on f , we get

$$\|\sigma_{2,k_{n_j}} - \sigma_{2,k_{n_{j'}}}\|_X \geq \frac{1}{L} \cdot 3^{N-2} = \frac{1}{L} \cdot r$$

for some constant $L > 0$ independent of ε . Indeed we can take L to be such that $L/2$ is the Lipschitz constant for f for distances larger than $\inf\{\|\sigma_{2,k_{n_j}} - \sigma_{2,k_{n_{j'}}}\|_X : j \neq j'\}$. Note that since f is uniformly continuous, $\inf\{\|\sigma_{2,k_{n_j}} - \sigma_{2,k_{n_{j'}}}\|_X : j \neq j'\}$ is strictly positive and independent of ε . In fact we have $\inf\{\|v_{J^{(k)}} - v_{J^{(k')}}\|_Y : k \neq k'\} \geq \theta = 3/4$ by the proof of item (1) in Lemma 3.0.1. So $\inf\{\|\sigma_{2,k_{n_j}} - \sigma_{2,k_{n_{j'}}}\|_X : j \neq j'\} \geq \eta$, where $\eta > 0$ is such that $\|\sigma - \sigma'\|_X < \eta$ implies $\|f(\sigma) - f(\sigma')\|_Y < 3/4$.

Combining this with the distances expressed in (2), and using property (β) on X , we must have for ε small enough,

$$\bar{\beta}_X \left(\frac{c_\infty^{\text{ATD}}(\tilde{\lambda})}{2L} \right) \leq \bar{\beta}_X \left(\frac{c_\infty^{\text{ATD}}(\tilde{\lambda})}{(1+3\varepsilon)L} \right) \leq \frac{83\varepsilon}{1+3\varepsilon}.$$

Since ε is arbitrary, we then get that

$$\bar{\beta}_X \left(\frac{c_\infty^{\text{ATD}}(\tilde{\lambda})}{2L} \right) = 0,$$

a contradiction. □

Remark 3.0.3. A superreflexive Banach space is one that has an equivalent norm which is uniformly convex (see for example [5, Appendix A]). A result of Bates, Johnson, Lindenstrauss, Preiss, and Schechtman [2] shows that a Banach space that is a uniform quotient of a superreflexive Banach space is itself superreflexive. Our result (by combining Theorem 2.0.1 and Theorem 3.0.2) gives a generalization of this for the case of separable Banach spaces. In fact, a uniformly convex Banach space has property (β) (see [29]); but there are separable spaces with property (β) which are not superreflexive (see for example [18]).

In [4] Baudier, Kalton and Lancien show that a Banach space which coarsely Lipschitz embeds into a separable Banach space with an equivalent norm with property (β) also has an equivalent norm with property (β) . Our result here gives the analog of this for uniform quotients.

4. QUANTITATIVE RESULTS

Recall the following asymptotic moduli.

Definition 4.0.4. Let X be an infinite-dimensional Banach space. Let $t \in (0, 1]$.

- (1) Modulus of asymptotic uniform convexity (AUC) [12, 15, 25]:

$$\bar{\delta}(t) = \inf_{\|x\|=1} \sup_{X_0} \inf_{\substack{z \in X_0 \\ \|z\| \geq t}} \{\|x + z\| - 1\},$$

where X_0 runs through all closed subspaces of X of finite codimension. The Banach space X is said to be asymptotically uniformly convex (AUC) if $\bar{\delta}(t) > 0$ for $t > 0$. A reflexive Banach space that is AUC is in particular called nearly uniformly convex (NUC).

- (2) Modulus of asymptotic uniform smoothness (AUS) [15, 25, 28]:

$$\bar{\rho}(t) = \sup_{\|x\|=1} \inf_{X_0} \sup_{\substack{z \in X_0 \\ \|z\| \leq t}} \{\|x + z\| - 1\},$$

where X_0 runs through all closed subspaces of X of finite codimension. The Banach space X is said to be asymptotically uniformly smooth (AUS) if $\lim_{t \rightarrow 0} \bar{\rho}(t)/t = 0$. A reflexive Banach space that is AUS is in particular called nearly uniformly smooth (NUS).

- (3) (β) -modulus [1]:

$$\bar{\beta}(t) = 1 - \sup \left\{ \inf \left\{ \frac{\|x + x_n\|}{2} : n \geq 1 \right\} \right\},$$

where the supremum is taken over all $x \in B_X$ and all sequences $(x_n)_{n \geq 1} \subseteq B_X$ with $\text{sep}((x_n)_{n \geq 1}) \geq t$. The Banach space X is said to have property (β) if $\bar{\beta}(t) > 0$ for $t > 0$.

Lemma 4.0.5. *For any infinite-dimensional Banach space X , and any $t \in (0, 1/2]$, we have $\bar{\beta}_X(t) \leq \bar{\delta}_X(2t)$.*

Proof. For $t \in (0, 1/2]$, denote $\beta = \bar{\beta}_X(t)$ and assume without loss of generality that $\beta > 0$. Let $x \in S_X$. Then for every sequence $(u_n)_{n \geq 1} \subseteq B_X$ with $\text{sep}((u_n)_{n \geq 1}) \geq t$, one has

$$\inf \left\{ \frac{\|x + u_n\|}{2} : n \geq 1 \right\} \leq 1 - \beta < \frac{1}{1 + \beta}.$$

Multiplying everything by $1 + \beta$ and taking the contrapositive, we get for any sequence $(x_n)_{n \geq 1}$ that if $\text{sep}((x_n)_{n \geq 1}) \geq (1 + \beta)t$, and if $\inf_{n \geq 1} \|(1 + \beta)x + x_n\| \geq 2$, then $\|x_n\| > 1 + \beta$ on a subsequence, and hence $\liminf \|x_n\| \geq 1 + \beta$.

Denote $T = 2t$. For a finite codimensional subspace X_0 , denote

$$\bar{\delta}_X(T, x, X_0) = \inf\{\|x + z\| - 1 : z \in X_0, \|z\| \geq T\};$$

and denote $\bar{\delta}_X(T, x)$ the supremum of $\bar{\delta}_X(T, x, X_0)$ over all finite codimensional subspaces X_0 .

Fix $\varepsilon > 0$. Let $x^* \in S_{X^*}$ be such that $x^*(x) = \|x\| = 1$. Denote by $X_1 = \ker(x^*)$. Let $z_1 \in X_1$ be such that $\|z_1\| \geq T$ and

$$\|x + z_1\| - 1 < \bar{\delta}_X(T, x, X_1) + \varepsilon \leq \bar{\delta}_X(T, x) + \varepsilon.$$

Let $z_1^* \in S_{X^*}$ be such that $z_1^*(z_1) = \|z_1\|$. Call $X_2 = X_1 \cap \ker(z_1^*)$, and let $z_2 \in X_2$ be such that $\|z_2\| \geq T$ and

$$\|x + z_2\| - 1 < \bar{\delta}_X(T, x, X_2) + \varepsilon \leq \bar{\delta}_X(T, x) + \varepsilon.$$

Continuing by induction, we find a sequence $(z_n)_{n \geq 1} \subseteq \ker(x^*)$ and functionals $(z_n^*)_{n \geq 1} \subseteq S_{X^*}$ such that for every $n \geq 1$ one has $z_n^*(z_n) = \|z_n\| \geq T$, and $z_{n+1} \in \bigcap_{i=1}^n \ker(z_i^*)$, and

$$\|x + z_n\| - 1 \leq \bar{\delta}_X(T, x) + \varepsilon.$$

Denote $x_n = x + z_n$. Then for $m > n$,

$$\|x_n - x_m\| = \|z_n - z_m\| \geq z_n^*(z_n - z_m) = z_n^*(z_n) = \|z_n\| \geq T.$$

So $\text{sep}((x_n)_{n \geq 1}) \geq T = 2t \geq (1 + \beta)t$. (Note that $\beta \leq 1$ by definition.) On the other hand, we have

$$\|(1 + \beta)x + x_n\| = \|(2 + \beta)x + z_n\| \geq x^*((2 + \beta)x + z_n) = 2 + \beta > 2.$$

As a result, we must have $\liminf \|x_n\| = \liminf \|x + z_n\| \geq 1 + \beta$.

This implies that $\beta = \bar{\beta}_X(t) \leq \bar{\delta}_X(2t, x) + \varepsilon$, and since $\varepsilon > 0$ and $x \in S_X$ are arbitrary, we get $\bar{\beta}_X(t) \leq \bar{\delta}_X(2t)$. □

Definition 4.0.6. We say that the (β) -modulus has power type p if there is a constant $C > 0$ so that $\bar{\beta}(t) \geq Ct^p$.

Proposition 4.0.7. *Let $1 < p < \infty$. There exist uniformly homeomorphic Banach spaces C_p and K_p so that C_p has (β) -modulus of power type p while K_p has no equivalent norm with (β) -modulus of power type p .*

Proof. Let $(G_n)_{n \geq 1}$ be a sequence of finite-dimensional Banach spaces dense in all finite-dimensional Banach spaces for the Banach-Mazur distance. Let $1 < p < \infty$. Johnson [14] introduced the space C_p given by $C_p = (\sum_{n=1}^{\infty} G_n)_{\ell_p}$. Kalton [17] considered the space $K_p = \left((\sum_{n=1}^{\infty} G_n)_{T_p} \right)^2$, where T_p is the p -convexification of Tsirelson space T (see for example [7]). Kalton proved in [17] that C_p and K_p are uniformly homeomorphic.

The computation in [9, Proposition 5.1] gives that C_p has (β) -modulus of power type p . On the other hand, Kalton remarks in [17] that K_p does not admit any

equivalent norm with AUC modulus of power type p . Assume that K_p admits an equivalent norm with (β) -modulus of power type p , that is $\bar{\beta}_{K_p}(t) \geq Ct^p$. By Lemma 4.0.5, we have $\bar{\beta}_{K_p}(t) \leq \bar{\delta}_{K_p}(2t)$. Thus for this norm we would obtain $\bar{\delta}_{K_p}(t) \geq C't^p$, which is in contradiction with Kalton's result. \square

Remark 4.0.8. In contrast with Theorems 2.0.1 and 3.0.2, as well as with [24, Proposition 4.1], the above proposition shows that we cannot compare $\bar{\beta}_{C_p}(t)$ and $\bar{\beta}_{K_p}(t)$ even under renorming, despite the fact that one is a uniform quotient of the other.

Furthermore, it was proved in [9] that if a Banach space Y is a uniform quotient of a Banach space Z , then for some constants C_1 and C_2 we have,

$$C_1 \bar{\beta}_Z(C_2 t) \leq \bar{\rho}_Y(t).$$

Taking $Z = C_p$ and $Y = K_p$, it follows from the above proposition that one can not replace the AUS modulus $\bar{\rho}_Y$ with the AUC modulus $\bar{\delta}_Y$ even under an equivalent norm on Y .

The rest of the article is devoted to an investigation of a renorming question. Recall that the AUS modulus is of power type p if there exists a constant C such that $\bar{\rho}(t) \leq Ct^p$. In [11] it is shown that if a separable Banach space X with AUS modulus of power type $p > 1$ is uniformly homeomorphic to another Banach space Y , then for any $q < p$, Y can be renormed to have AUS modulus of power type q . It is natural to ask whether the same holds for the (β) -modulus, namely:

Assume that two Banach spaces are uniformly homeomorphic and that one of them has (β) -modulus of power type p . Given an arbitrary $q > p$, does the other space always have an equivalent norm with (β) -modulus of power type q ?

In the case of the counterexample provided by the pair of spaces C_p and K_p , where $1 < p < \infty$, we can give a positive answer to the above question.

Proposition 4.0.9. *Assume that X is a separable reflexive Banach space which is uniformly homeomorphic to a space with $\bar{\rho}(t) \leq C_1 t^p$, $1 < p < \infty$, and that the dual space X^* is uniformly homeomorphic to a space with $\bar{\rho}(t) \leq C_2 t^{p'}$ where $1/p + 1/p' = 1$. Then for any $\gamma > 0$, X admits an equivalent norm with $\bar{\beta}(t) \geq Ct^{p+\gamma}$.*

Proof. By the result in [11], the space X has an equivalent norm $\|\cdot\|_1$ with $\bar{\rho}_{\|\cdot\|_1}(t)$ of power type $p - \varepsilon$. Similarly, for any $\varepsilon_1 > 0$, X^* has an equivalent norm $\|\cdot\|_2^*$ with $\bar{\rho}_{\|\cdot\|_2^*}(t)$ of power type $p' - \varepsilon_1$. By duality [28] (see also [11]), X has $\bar{\delta}_{\|\cdot\|_2}(t)$ of power type $p + \varepsilon$ provided we choose ε_1 accordingly. Then by [15, Proposition 2.10] (see also [26]), X admits an equivalent norm $\|\cdot\|$ with both $\bar{\rho}(t)$ of power type $p - \varepsilon$ and $\bar{\delta}(t)$ of power type $p + \varepsilon$.

It was shown in [19] that if a norm is both NUC and NUS then it also has property (β) . Following the same avenue, it was proved in [10] that a norm with the above power types for $\bar{\rho}(t)$ and $\bar{\delta}(t)$ has a (β) -modulus $\bar{\beta}(t)$ of power type

$$\frac{(p - \varepsilon)(p + \varepsilon) - (p - \varepsilon)}{(p - \varepsilon) - 1} = p + o(\varepsilon).$$

\square

Corollary 4.0.10. *For every $\varepsilon > 0$ the space $K_p = \left((\sum_{n=1}^{\infty} G_n)_{T_p} \right)^2$, $1 < p < \infty$, has an equivalent norm with (β) -modulus of power type $p + \varepsilon$.*

Proof. By [17], K_p is uniformly homeomorphic to C_p and $\bar{\rho}_{C_p}(t) \leq C_1 t^p$. On the other hand, we have $K_p^* = \left((\sum_{n=1}^{\infty} G_n^*)_{T_p^*} \right)^2 = \left((\sum_{n=1}^{\infty} G_n^*)_{T_{p'}} \right)^2$. Since $(G_n^*)_{n \geq 1}$ is also dense in Banach-Mazur metric, we obtain by the same theorem in [17] that $\left((\sum_{n=1}^{\infty} G_n^*)_{T_{p'}} \right)^2$ is uniformly homeomorphic to $C_{p'}$; and $\bar{\rho}_{C_{p'}}(t) \leq C_2 t^{p'}$. \square

5. ACKNOWLEDGMENTS

This work was initiated while the authors attended the NSF funded Workshop in Linear Analysis and Probability at Texas A&M University during Summer 2012, and was finalized while the first two authors attended this same Workshop during Summer 2013.

The authors would like to thank the anonymous referee for his/her invaluable comments that helped smooth out the presentation of this paper.

REFERENCES

- [1] J. M. Ayerbe, T. Domínguez Benavides, and S. Francisco Cutillas, *Some noncompact convexity moduli for the property (β) of Rolewicz*, Comm. Appl. Nonlinear Anal. **1** (1994), no. 1, 87–98.
- [2] S. Bates, W. B. Johnson, J. Lindenstrauss, D. Preiss, and G. Schechtman, *Affine approximation of Lipschitz functions and nonlinear quotients*, Geom. Funct. Anal. **9** (1999), no. 6, 1092–1127.
- [3] Florent Baudier, *Metrical characterization of super-reflexivity and linear type of Banach spaces*, Arch. Math. **89** (2007), no. 5, 419–429.
- [4] F. Baudier, N. J. Kalton, and G. Lancien, *A new metric invariant for Banach spaces*, Studia Math. **199** (2010), no. 1, 73–94.
- [5] Yoav Benyamini and Joram Lindenstrauss, *Geometric nonlinear functional analysis. Vol. 1*, American Mathematical Society Colloquium Publications, vol. 48, American Mathematical Society, Providence, RI, 2000.
- [6] J. Bourgain, *The metrical interpretation of superreflexivity in Banach spaces*, Israel J. Math. **56** (1986), no. 2, 222–230.
- [7] Peter G. Casazza and Thaddeus J. Shura, *Tsirel'son's space*, Lecture Notes in Mathematics, vol. 1363, Springer-Verlag, Berlin, 1989. With an appendix by J. Baker, O. Slotterbeck and R. Aron.
- [8] Jeff Cheeger and Bruce Kleiner, *Realization of metric spaces as inverse limits, and bilipschitz embedding in L_1* , Geom. Funct. Anal. **23** (2013), no. 1, 96–133.
- [9] S. J. Dilworth, Denka Kutzarova, G. Lancien, and N. L. Randrianarivony, *Asymptotic geometry of Banach spaces and uniform quotient maps*, Proc. Amer. Math. Soc. **142** (2014), no. 8, 2747–2762.
- [10] S. J. Dilworth, Denka Kutzarova, N. Lovasoa Randrianarivony, J. P. Revalski, and N. V. Zhivkov, *Compactly uniformly convex spaces and property (β) of Rolewicz*, J. Math. Anal. Appl. **402** (2013), no. 1, 297–307.
- [11] G. Godefroy, N. J. Kalton, and G. Lancien, *Szlenk indices and uniform homeomorphisms*, Trans. Amer. Math. Soc. **353** (2001), no. 10, 3895–3918 (electronic).
- [12] R. Huff, *Banach spaces which are nearly uniformly convex*, Rocky Mountain J. Math. **10** (1980), no. 4, 743–749.
- [13] Robert C. James, *Weak compactness and reflexivity*, Israel J. Math. **2** (1964), 101–119.
- [14] William B. Johnson, *Factoring compact operators*, Israel J. Math. **9** (1971), 337–345.
- [15] William B. Johnson, Joram Lindenstrauss, David Preiss, and Gideon Schechtman, *Almost Fréchet differentiability of Lipschitz mappings between infinite-dimensional Banach spaces*, Proc. London Math. Soc. (3) **84** (2002), no. 3, 711–746.

- [16] William B. Johnson and Gideon Schechtman, *Diamond graphs and super-reflexivity*, J. Topol. Anal. **1** (2009), no. 2, 177–189.
- [17] N. J. Kalton, *Examples of uniformly homeomorphic Banach spaces*, Israel J. Math. **194** (2013), no. 1, 151–182.
- [18] D. N. Kutzarova, *On condition (β) and Δ -uniform convexity*, C. R. Acad. Bulgare Sci. **42** (1989), no. 1, 15–18.
- [19] Denka Kutzarova, *An isomorphic characterization of property (β) of Rolewicz*, Note Mat. **10** (1990), no. 2, 347–354.
- [20] ———, *k - β and k -nearly uniformly convex Banach spaces*, J. Math. Anal. Appl. **162** (1991), no. 2, 322–338.
- [21] T. J. Laakso, *Ahlfors Q -regular spaces with arbitrary $Q > 1$ admitting weak Poincaré inequality*, Geom. Funct. Anal. **10** (2000), no. 1, 111–123.
- [22] Tomi J. Laakso, *Plane with A_∞ -weighted metric not bi-Lipschitz embeddable to \mathbb{R}^n* , Bull. London Math. Soc. **34** (2002), no. 6, 667–676.
- [23] Vegard Lima and N. Lovasoa Randrianarivony, *Property (β) and uniform quotient maps*, Israel J. Math. **192** (2012), no. 1, 311–323.
- [24] Manor Mendel and Assaf Naor, *Markov convexity and local rigidity of distorted metrics*, J. Eur. Math. Soc. (JEMS) **15** (2013), no. 1, 287–337.
- [25] V. D. Milman, *Geometric theory of Banach spaces. II. Geometry of the unit ball*, Uspehi Mat. Nauk **26** (1971), 73–149 (Russian). English translation: Russian Math. Surveys **26** (1971), 79–163.
- [26] Edward W. Odell and Thomas Schlumprecht, *Embeddings into Banach spaces with finite dimensional decompositions*, RACSAM. Rev. R. Acad. Cienc. Ser. A Mat. **100** (2006), no. 1-2, 295–323.
- [27] Mikhail Ostrovskii, *On metric characterizations of the Radon-Nikodým and related properties of Banach spaces*, preprint.
- [28] Stanisław Prus, *Nearly uniformly smooth Banach spaces*, Boll. Un. Mat. Ital. B **7** (1989), no. 3, 507–521.
- [29] S. Rolewicz, *On drop property*, Studia Math. **85** (1986), no. 1, 27–35 (1987).
- [30] ———, *On Δ -uniform convexity and drop property*, Studia Math. **87** (1987), no. 2, 181–191.
- [31] Jeremy T. Tyson, *Bi-Lipschitz embeddings of hyperspaces of compact sets*, Fund. Math. **187** (2005), no. 3, 229–254.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTH CAROLINA, COLUMBIA, SC 29208, USA.

E-mail address: dilworth@math.sc.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN, URBANA, IL 61801, USA.

E-mail address: denka@math.uiuc.edu

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, SAINT LOUIS UNIVERSITY, ST. LOUIS, MO 63103, USA.

E-mail address: nrandria@slu.edu